

WHO CLASSIFICATION OF ENDOCRINE TUMOURS Full Version

WHO classification of endocrine tumours is a pivotal framework used by pathologists, oncologists, and medical professionals worldwide to categorize and understand the diverse spectrum of endocrine neoplasms. This classification system is continually updated by the World Health Organization (WHO) to incorporate new scientific insights, improve diagnostic accuracy, and guide effective treatment strategies. Endocrine tumours originate from hormone-producing glands, including the thyroid, parathyroid, adrenal glands, and neuroendocrine cells dispersed throughout various organs. Proper classification is essential for prognosis determination, therapeutic planning, and facilitating research efforts. In this comprehensive overview, we explore the WHO classification of endocrine tumours, detailing the categories, key features, diagnostic criteria, and latest updates to this vital system.

Understanding the WHO Classification of Endocrine Tumours

The WHO classification of endocrine tumours is a histopathological and molecular framework that categorizes tumors based on their origin, cellular differentiation, malignancy potential, and molecular features. The system emphasizes a multidisciplinary approach, combining morphological assessment with immunohistochemistry, molecular genetics, and clinical data.

Key objectives of the WHO classification include:

- Standardizing diagnosis across institutions and countries
- Facilitating research and clinical trials
- Improving patient management and prognosis prediction
- Incorporating advances in molecular pathology

The classification is periodically revised, with the latest edition published in 2022, reflecting significant advances in understanding endocrine tumours.

Major Categories in the WHO Classification of Endocrine Tumours

The classification broadly divides endocrine tumours into several main categories, each with specific subtypes, based on tissue of origin and biological behavior.

1. Tumours of the Thyroid Gland

Thyroid tumours are among the most common endocrine neoplasms. They are classified into benign and malignant categories.

Benign Tumours:

- Follicular adenoma
- Hürthle cell adenoma

Malignant Tumours:

- Papillary thyroid carcinoma (PTC)
- Follicular thyroid carcinoma (FTC)
- Medullary thyroid carcinoma (MTC)
- Anaplastic thyroid carcinoma (ATC)

Key features:

- Differentiation status
- Presence of specific genetic mutations (e.g., BRAF, RAS)
- Histological architecture

2. Parathyroid Tumours

Parathyroid tumours are predominantly benign but can occasionally be malignant.

Main types:

- Parathyroid adenoma (most common)
- Parathyroid hyperplasia
- Parathyroid carcinoma (rare but aggressive)

Diagnostic highlights:

- Elevated serum calcium and parathyroid hormone levels
- Histopathological features distinguishing benign from malignant

3. Adrenal Gland Tumours

Adrenal tumours include those arising from the cortex and medulla, with distinct classifications.

Adrenal Cortical Tumours:

- Adenomas (benign)
- Carcinomas (malignant)
- Myelolipomas and other rare tumours

Adrenal Medullary Tumours:

- Pheochromocytomas
- Paragangliomas (extra-adrenal sympathetic or parasympathetic)

Features:

- Hormone secretion profiles
- Malignancy potential based on invasion and metastasis
- Molecular markers (e.g., SDH mutations)

4. Neuroendocrine Tumours (NETs)

Neuroendocrine tumours are a diverse group originating from neuroendocrine cells scattered throughout the body.

Primary sites include:

- Gastroenteropancreatic (GEP) system
- Lung
- Thymus
- Other organs

Classification:

- Well-differentiated neuroendocrine tumours (NETs)

- Poorly differentiated neuroendocrine carcinomas (NECs)

Grading and staging:

- Based on mitotic count and Ki-67 proliferation index
- Graded as G1, G2, or G3

Key features:

- Hormone secretion (functional vs. non-functional)
- Morphological subtype (e.g., carcinoid, small cell carcinoma)

Histopathological and Molecular Features in WHO Classification

The latest WHO classification emphasizes integrating histopathology with molecular diagnostics. Key molecular markers and genetic alterations are now integral to tumor classification.

Common molecular features include:

- BRAF, RAS, RET mutations in thyroid carcinomas
- MEN1, DAXX, ATRX mutations in pancreatic NETs
- SDH mutations in paragangliomas and pheochromocytomas
- TP53 mutations in anaplastic carcinomas

This integration enhances diagnostic precision, allows for targeted therapies, and provides prognostic information.

Update Highlights from the 2022 WHO Classification

The 2022 edition introduced several notable updates:

- Refinement of grading criteria for neuroendocrine tumours, emphasizing proliferation markers
- Clarification of molecular subtypes within thyroid carcinomas, especially papillary and follicular types
- Recognition of new entities, such as well-differentiated grade 3 neuroendocrine tumours
- Inclusion of emerging molecular data supporting targeted therapies
- Standardization of terminology to reduce ambiguity in diagnosis

These updates reflect ongoing research and technological advances in molecular pathology.

Practical Implications of WHO Classification in Clinical Practice

Accurate classification per WHO standards influences every aspect of patient care:

Diagnosis:

- Guides biopsy interpretation
- Determines the need for molecular testing

Treatment planning:

- Surgical approach
- Use of targeted therapies
- Surveillance strategies

Prognosis estimation:

- Differentiation status
- Tumor grade and stage
- Molecular features

Research and clinical trials:

- Stratification of patients
- Development of novel therapies

Conclusion

The WHO classification of endocrine tumours is a comprehensive, dynamic system that plays a crucial role in the diagnosis, management, and research of endocrine neoplasms. Its integration of histopathology with molecular genetics has revolutionized understanding and treatment of these tumours. Staying updated with the latest WHO classifications ensures clinicians and pathologists can provide accurate diagnoses and optimal care, ultimately improving patient outcomes. As advancements continue, the classification will evolve further, incorporating new biomarkers and targeted therapies, reaffirming its importance in the field of endocrine oncology.

WHO classification of endocrine tumours

The WHO classification of endocrine tumours represents a critical framework used worldwide by pathologists, oncologists, and researchers to diagnose, categorize, and manage tumors originating from endocrine glands. These tumors encompass a diverse group of neoplasms arising from endocrine tissues such as the thyroid, parathyroid, adrenal glands, and neuroendocrine cells throughout the body. Over the years, this classification has evolved significantly, integrating advances in histopathology, molecular genetics, and clinical behavior to provide a comprehensive and standardized approach. Its importance lies not only in improving diagnostic accuracy but also in guiding treatment strategies and prognostication.

Historical Context and Development of WHO Classification

Understanding the evolution of the WHO classification helps appreciate its current utility. Initially, endocrine tumors were classified based solely on histological appearance and clinical features. However, as molecular biology advanced, it became evident that tumors with similar histology could have vastly different genetic profiles and behaviors. The first WHO classifications for endocrine tumors were published in the 1980s, with subsequent updates in 1992, 2004, 2010, and most recently in 2017.

Key milestones include:

- Integration of molecular markers to refine tumor subtypes.
- Emphasis on grading and staging systems to assess malignancy.
- Inclusion of new entities based on genetic alterations.

The latest WHO classification (2017) emphasizes a multidisciplinary approach, combining morphology, immunohistochemistry, and molecular findings.

Core Principles and Structure of the WHO Classification

The WHO classification is designed to be comprehensive yet adaptable, categorizing endocrine tumors based on:

- Histological features: cell morphology, architecture, mitotic activity.
- Immunohistochemical profiles: hormone production, neuroendocrine markers.
- Molecular genetics: presence of specific mutations or gene alterations.
- Behavioral criteria: benign, uncertain malignant potential, or malignant.

The classification groups endocrine tumors mainly into:

- Tumors of the thyroid gland
- Tumors of the parathyroid glands
- Tumors of the adrenal cortex and medulla
- Neuroendocrine tumors (NETs) of other sites

Each category includes specific tumor types with well-defined criteria, facilitating consistent diagnosis across institutions.

Thyroid Tumors

Thyroid neoplasms are among the most common endocrine tumors, and their classification is vital for prognosis and management.

Major Categories

- Differentiated thyroid carcinomas (DTCs)
 - Papillary thyroid carcinoma (PTC)
 - Follicular thyroid carcinoma (FTC)
 - Hurthle cell carcinoma
- Medullary thyroid carcinoma (MTC)
- Anaplastic thyroid carcinoma
- Thyroid adenomas

Features and Features of Each Type

- Papillary thyroid carcinoma:
 - Most common thyroid cancer (about 80%)
 - Characteristic nuclear features (orphan Annie eyes, nuclear grooves)
 - Often associated with BRAF mutations
 - Good prognosis with appropriate treatment

- Follicular thyroid carcinoma:
 - Accounts for ~10-15% of thyroid cancers
 - Distinguished from adenoma by capsular and vascular invasion
 - Associated with RAS mutations and PAX8-PPAR γ rearrangements

- Medullary thyroid carcinoma:
 - Arises from parafollicular C cells
 - Produces calcitonin
 - Can be sporadic or familial (MEN 2 syndromes)
 - Associated with RET mutations

- Anaplastic thyroid carcinoma:
 - Highly aggressive and undifferentiated
 - Poor prognosis
 - Often arises from pre-existing differentiated carcinomas

Pros and Cons of WHO Classification for Thyroid Tumors

| Pros | Cons |

|-----|-----|

- | Clear histological and molecular criteria improve accuracy | Some tumors may have overlapping features, complicating diagnosis |

- | Guides targeted therapies (e.g., RET inhibitors for MTC) | Rare variants may not fit neatly into categories |

- | Facilitates prognostication and management planning | Requires specialized expertise and resources |

Parathyroid Tumors

Parathyroid neoplasms are relatively rare but clinically significant due to their role in primary hyperparathyroidism.

Types and Features

- Adenoma (most common)
- Hyperplasia
- Carcinoma (rare)

Features

- Parathyroid adenomas:
 - Usually solitary
 - Composed of chief cells, with a thin capsule
 - Typically benign with excellent prognosis
- Parathyroid carcinoma:
 - Larger, invasive, may invade surrounding tissues
 - Histologically challenging; criteria include capsular, vascular, or perineural invasion
 - Rare but aggressive

Strengths and Limitations

- The WHO classification emphasizes invasion and metastasis for malignancy diagnosis, which can sometimes be subtle histologically.
- Accurate diagnosis influences surgical management and follow-up.

Adrenal Tumors

Adrenal tumors are diverse, including cortical, medullary, and metastatic lesions.

Adrenal Cortical Tumors

- Adenomas (benign)
- Carcinomas (malignant)
- Myelolipomas and other benign lesions

Features

- Adrenal cortical adenomas:
 - Usually nonfunctioning or secreting cortisol, aldosterone, or androgens

- Well-circumscribed, benign features
- Adrenal cortical carcinomas:
 - Larger, irregular, invasive
 - Mitotic figures, nuclear atypia, necrosis, and invasion are hallmarks
 - Weiss criteria are used for diagnosis

Adrenal Medullary Tumors

- Pheochromocytomas:
 - Derived from chromaffin cells
 - Secrete catecholamines
 - Classified as benign or malignant based on invasion and metastasis

Features and Prognostic Indicators

- Malignancy in adrenal tumors is confirmed by invasion or metastasis, not solely histology.
- Molecular markers and scoring systems (e.g., PASS) assist in risk stratification.

Neuroendocrine Tumors (NETs) of Other Sites

NETs are characterized by neuroendocrine differentiation and can occur throughout the body.

Classification and Features

- Gastroenteropancreatic neuroendocrine tumors (GEP-NETs)
- Well-differentiated tumors (NETs) and poorly differentiated neuroendocrine carcinomas (NECs)
- Graded based on mitotic count and Ki-67 index into G1, G2, G3
- Pulmonary NETs
 - Typical carcinoid
 - Atypical carcinoid
 - Large cell neuroendocrine carcinoma
 - Small cell lung carcinoma

Advantages of WHO Classification

- Incorporates proliferation indices (mitotic rate, Ki-67) for grading, which correlates with prognosis.
- Recognizes distinct biological behaviors and treatment responses based on grade and differentiation.

Limitations

- Requires precise histological and immunohistochemical assessment.
- Some tumors demonstrate heterogeneity, complicating classification.

Molecular Advances and Future Directions

The WHO classification increasingly integrates molecular data, recognizing that genetic alterations can refine diagnosis and predict behavior.

- Mutations such as BRAF, RAS, RET, and TP53 are incorporated into diagnostic criteria.
- Gene expression profiles aid in subclassification, especially for poorly differentiated tumors.
- Targeted therapies are emerging based on molecular alterations, making these classifications clinically relevant.

The future of the WHO classification is likely to involve:

- More precise molecular subclassifications
- Personalized risk assessments
- Incorporation of emerging biomarkers

Conclusion

The WHO classification of endocrine tumours remains a cornerstone in endocrine pathology, providing a structured, evidence-based approach to tumor diagnosis and management. Its comprehensive nature?combining histological, immunohistochemical, and molecular features?enhances diagnostic precision, facilitates prognostication, and guides therapy. While it has notable strengths, including standardization and integration of advances, challenges such as evolving molecular data and tumor heterogeneity persist. Continued refinement, incorporating new molecular insights, will ensure that the WHO classification remains relevant and invaluable in the era of personalized medicine.

In summary, understanding the WHO classification of endocrine tumors is essential for clinicians

and pathologists alike, enabling consistent diagnosis, better patient outcomes, and the advancement of research in endocrine oncology.

Question What is the WHO classification of endocrine tumours based on? The WHO classification of endocrine tumours is based on histopathological features, cellular origin, differentiation status, and molecular characteristics to categorize tumours into benign, uncertain malignant potential, and malignant types. How does the WHO classification differentiate between benign and malignant endocrine tumours? The WHO classification uses criteria such as cellular atypia, mitotic activity, invasion, and metastasis presence to distinguish benign from malignant endocrine tumours, with specific categories like neuroendocrine tumors, carcinomas, and neoplasms of uncertain malignant potential. What are the key updates in the latest WHO classification of endocrine tumours? The latest updates incorporate molecular profiling, grading systems based on proliferation indices (like Ki-67), and refined criteria for borderline or uncertain malignant potential tumours to improve diagnostic accuracy and prognostication. How does the WHO classification impact clinical management of endocrine tumours? It guides treatment decisions by providing a standardized framework for diagnosis, grading, and staging, helping clinicians determine prognosis and appropriate therapeutic strategies for patients with endocrine tumours. Which endocrine tumours are most commonly classified under the WHO system? Commonly classified tumours include thyroid carcinomas (papillary, follicular, medullary, anaplastic), neuroendocrine tumours of the pancreas, adrenal cortical tumours, and parathyroid neoplasms. What role does molecular pathology play in the WHO classification of endocrine tumours? Molecular pathology aids in identifying genetic mutations and molecular subtypes, which refine classification, improve diagnostic precision, and can influence targeted therapy options in endocrine tumours.

Related keywords: endocrine tumors, WHO tumor classification, neuroendocrine tumors, endocrine neoplasms, tumor grading, tumor staging, endocrine pathology, tumor histology, neuroendocrine carcinoma, tumor biomarkers

Who classification of tumours of the urinary syst

WHO Classification of Tumours of the Urinary System

The World Health Organization (WHO) classification of tumours of the urinary system provides a comprehensive framework for diagnosing, categorizing, and understanding the diverse neoplasms that affect the kidneys, urinary bladder, ureters, and other related structures. This classification aids pathologists, urologists, oncologists, and researchers in standardizing diagnoses, informing treatment strategies, and facilitating epidemiological studies. The WHO classification is regularly updated to reflect advances in molecular pathology, histopathology, and clinical research, ensuring

it remains a vital reference in contemporary medicine.

Overview of the WHO Classification of Tumours of the Urinary System

The WHO classification groups urinary system tumors based on their histogenesis, morphological features, molecular characteristics, and clinical behavior. It covers tumors of the kidneys, renal pelvis, ureters, urinary bladder, and urethra. Each tumor type is assigned a specific code, description, and grading/staging parameters that assist clinicians in prognosis and management.

Key features of the classification include:

- Categorization into benign and malignant neoplasms
- Recognition of precursor lesions
- Incorporation of molecular markers
- Grading systems that reflect tumor aggressiveness

Renal Tumours

Renal tumors constitute a significant portion of urinary system neoplasms. The most common malignant renal tumor is renal cell carcinoma (RCC), but several other benign and malignant tumors also occur.

1. Renal Cell Carcinoma (RCC)

Renal cell carcinoma accounts for approximately 85% of malignant kidney tumors in adults. The WHO classifies RCC into various subtypes based on histological and molecular features:

- **Clear Cell RCC** (most common subtype, ~70%)
- **Papillary RCC** (~10-15%)
- **Chromophobe RCC** (~5%)
- **Collecting Duct Carcinoma** (Bellini duct carcinoma)
- **MiT family translocation RCC**

Histological and Molecular Features:

- Clear cell RCC shows clear cytoplasm with a rich vascular network.
- Papillary RCC exhibits papillary architecture with foamy macrophages.

- Chromophobe RCC displays pale cytoplasm with prominent cell membranes.
- Molecular markers such as VHL gene mutations are characteristic of clear cell RCC.

Grading and Staging:

- Fuhrman grading system assesses nuclear features.
- TNM staging evaluates tumor size, node involvement, and metastasis.

2. Renal Tumors of Uncertain Malignant Potential (UMP)

Includes lesions like:

- Oncocytoma (benign)
- Hybrid tumors with features of oncocytoma and chromophobe RCC

3. Benign Renal Tumors

- Oncocytoma: Composed of oncocytic cells with granular eosinophilic cytoplasm.
- Angiomyolipoma: Composed of blood vessels, smooth muscle, and fat; associated with tuberous sclerosis.

Urothelial (Transitional Cell) Tumours

Urothelial tumors are the most common neoplasms of the urinary bladder and urothelium lining the renal pelvis and ureters.

1. Non-invasive Urothelial Carcinoma

- Papillary Urothelial Neoplasm of Low Malignant Potential (PUNLMP): Minimal cytological atypia.
- Non-invasive papillary carcinoma (Ta): Confined to the mucosa, with papillary growth.
- Carcinoma in situ (CIS): Flat, high-grade lesion with a high risk of invasion.

2. Invasive Urothelial Carcinoma

- Muscle-invasive carcinoma (T2 and beyond): Penetrates muscularis propria.
- Variants and special types: Include micropapillary, nested, sarcomatoid, and plasmacytoid

variants, which often have worse prognosis.

Histological Grading:

- WHO/ISUP grading system classifies tumors into low-grade and high-grade based on nuclear features and architecture.

Molecular Features:

- FGFR3 mutations are common in low-grade tumors.
- TP53 mutations often associated with high-grade, invasive tumors.

Other Malignant Tumours of the Urinary System

Besides RCC and urothelial carcinomas, other less common malignant tumors include:

1. Wilms Tumor (Nephroblastoma)

Primarily affects children; characterized by triphasic histology (blastemal, epithelial, stromal).

2. Sarcomas

- Leiomyosarcoma: Arises from smooth muscle.
- Angiosarcoma: Malignant vascular tumor.

3. Lymphomas and Metastases

- Lymphomatous infiltration can involve the kidney.
- Metastatic tumors from lung, breast, or gastrointestinal tract.

Benign Tumours of the Urinary System

Benign neoplasms are less frequent but include:

1. Oncocytoma

- Composed of oncocytic cells with abundant mitochondria.
- Usually asymptomatic and discovered incidentally.

2. Urothelial Papillomas

- Small, papillary lesions with benign features.
- Rarely progress to malignancy but require follow-up.

3. Angiomyolipoma

- Common benign tumor of the kidney, especially in tuberous sclerosis.
- Composed of blood vessels, smooth muscle, and fat.

Molecular Pathology and Diagnostic Advances

The WHO classification increasingly incorporates molecular diagnostics, which:

- Aid in differentiating subtypes
- Predict prognosis
- Guide targeted therapies

Examples include:

- VHL gene mutations in clear cell RCC
- FGFR3 mutations in low-grade urothelial tumors
- Chromosomal aberrations and gene expression profiles

Prognostic Factors and Clinical Implications

Understanding tumor classification influences prognosis and treatment:

- Tumor subtype and grade determine aggressiveness.
- Molecular features inform targeted therapy options.
- Accurate staging guides surgical and systemic management.

Prognostic Indicators:

- Histological subtype
- Grade and stage
- Molecular markers
- Presence of invasion or metastasis

Conclusion

The WHO classification of tumours of the urinary system provides a detailed, standardized approach to diagnosing and understanding these diverse neoplasms. Recognizing the different tumor types?ranging from benign to highly malignant?helps clinicians tailor treatment, predict outcomes, and advance research. With ongoing discoveries in molecular pathology, this classification continues to evolve, enhancing precision medicine in urology and oncology.

Key Takeaways:

- The classification covers renal tumors, urothelial carcinomas, and rare neoplasms.
- Histological, molecular, and clinical features are essential for accurate diagnosis.
- Tumor grading and staging are crucial for management decisions.
- Advances in molecular pathology are increasingly integrated into classification systems.

For healthcare professionals, staying updated with the latest WHO classifications ensures accurate diagnosis and optimal patient care in the complex landscape of urinary system tumors.

Who Classification of Tumours of the Urinary System: A Comprehensive Guide

Understanding the who classification of tumours of the urinary system is essential for clinicians, pathologists, and medical students alike. This classification system provides a standardized framework that helps in diagnosing, researching, and treating various neoplasms affecting the urinary tract. From kidneys to ureters and bladder, these tumours exhibit diverse histological features and clinical behaviors. The WHO classification serves as a critical tool in guiding management strategies and prognostic assessments.

Introduction to the WHO Classification of Urinary System Tumours

The World Health Organization (WHO) periodically updates its classification of tumours of the urinary system to reflect advances in histopathology, molecular biology, and clinical research. The latest editions categorize tumours based on their tissue of origin, cellular morphology, and biological behavior. This systematic approach enhances diagnostic accuracy and facilitates communication among healthcare providers.

The WHO classification of tumours of the urinary system encompasses tumours arising from:

- Kidney (renal cell carcinomas and others)
- Renal pelvis and ureter
- Urinary bladder
- Urethra

Each site has distinct tumour types with specific histological and molecular features, which are crucial for prognosis and therapy.

Significance of the WHO Classification

- Standardization: Provides a uniform language for diagnosis.
- Prognostic Value: Helps predict disease course.
- Therapeutic Guidance: Influences treatment choices.
- Research Framework: Facilitates clinical and translational research.

Tumours of the Kidney

The kidney is the most common site for urinary system tumours, with renal cell carcinoma (RCC) being the predominant malignant tumour.

Main Categories of Renal Tumours

- Renal Cell Carcinomas (RCCs)
 - The most common malignant kidney tumours in adults.
 - Subtypes based on histology and molecular features:
 - Clear cell RCC
 - Papillary RCC
 - Chromophobe RCC
 - Collecting duct carcinoma
 - MiT family translocation RCCs
- Benign Renal Tumours

- Oncocytoma
- Angiomyolipoma
- Renal cysts (non-neoplastic but important in differential diagnosis)

Key Features of RCC Subtypes

- Clear cell RCC: Characterized by cells with clear cytoplasm, often associated with VHL gene mutations.
- Papillary RCC: Exhibits papillary architecture, often with gains of chromosomes 7 and 17.
- Chromophobe RCC: Has pale eosinophilic cytoplasm, with distinct cell membranes, often associated with BHD syndrome.

Tumours of the Renal Pelvis and Ureter

These tumours are predominantly transitional cell carcinomas (urothelial carcinomas).

Main Tumour Types

- Urothelial carcinoma: The most common, similar to bladder tumours.
- Squamous cell carcinoma: Rare, often linked to chronic inflammation.
- Adenocarcinoma: Very rare.

Pathogenesis and Features

- Arise from the transitional epithelium lining the renal pelvis and ureter.
- Often associated with urinary tract stones and chronic irritation.

Tumours of the Urinary Bladder

Bladder tumours are among the most common urinary system neoplasms.

Major Categories

- Urothelial (transitional cell) carcinoma: The majority.
- Squamous cell carcinoma: Usually linked to schistosomiasis or chronic irritation.
- Adenocarcinoma: Rare, may arise from urachal remnants.
- Small cell carcinoma: Neuroendocrine differentiation.

Grading and Staging

- Graded based on cellular atypia and architectural disorder.
- Staged according to tumor invasion depth and spread.

Key Subtypes of Urothelial Carcinoma

- Papillary urothelial carcinoma: Often low-grade, papillary architecture.
- Flat carcinoma in situ: High-grade, flat lesions with potential for invasive disease.
- Invasive urothelial carcinoma: Penetrates muscularis propria or beyond.

Tumours of the Urethra

Urethral cancers are rare but can include:

- Urothelial carcinoma (most common)
- Squamous cell carcinoma
- Adenocarcinoma

Their classification relies on similar histopathological features as bladder tumours.

Molecular and Histological Classification

The WHO classification of tumours of the urinary system integrates molecular markers with histology to refine diagnosis:

- Clear cell RCC: VHL gene mutations, VEGF pathway alterations.
- Papillary RCC: MET gene mutations.
- Chromophobe RCC: Chromosomal losses, mitochondrial abnormalities.
- Urothelial carcinoma: FGFR3 mutations, TP53 alterations.

Incorporating molecular data improves prognostication and opens avenues for targeted therapies.

Practical Approach to Classification

Step 1: Histopathological Examination

- Identify tumour origin and morphology.
- Determine histological subtype.

Step 2: Tumour Grade and Stage

- Use WHO grading systems.
- Assess invasion extent and spread.

Step 3: Molecular Studies

- When indicated, perform genetic testing for specific mutations.

Summary Table: Key Tumour Types in the WHO Classification

Site	Main Tumour Types	Notable Features
Kidney	Renal cell carcinoma (clear cell, papillary, chromophobe), oncocytoma	Histology, molecular profile
Renal Pelvis/Ureter	Urothelial carcinoma, squamous cell carcinoma	Association with stones/inflammation
Urinary Bladder	Urothelial carcinoma (papillary, flat, invasive), squamous cell, adenocarcinoma	Grading, staging, molecular markers
Urethra	Urothelial, squamous, adenocarcinoma	Rarity necessitates careful classification

Conclusion

The WHO classification of tumours of the urinary system provides a detailed, histology-based framework that guides diagnosis, prognosis, and treatment. Recognizing the distinct types of tumours, their molecular underpinnings, and their clinical behaviors is fundamental to effective management. As research advances, this classification continues to evolve, integrating genomic data to refine our understanding of urinary tract neoplasms.

Final Thoughts

Staying updated with the latest WHO classifications is vital for healthcare professionals involved in diagnosing and treating urinary system tumours. A thorough understanding of the diverse tumour types, their pathological features, and molecular characteristics enables personalized patient care and contributes to improved outcomes.

Question What is the WHO classification of tumors of the urinary system based on? The WHO classification is based on histopathological features, tumor morphology, grade, and molecular characteristics to categorize tumors of the urinary system accurately. How are urothelial carcinomas categorized in the WHO classification? Urothelial carcinomas are classified into non-invasive papillary carcinomas, carcinoma in situ, and invasive carcinomas with various histological subtypes and grades. What are the main categories of renal tumors according to the WHO classification? Main categories include clear cell renal cell carcinoma, papillary renal cell carcinoma, chromophobe renal cell carcinoma, and other rare types like oncocytoma and collecting duct carcinoma. How does the WHO classify tumors of the prostate and bladder? Prostate tumors are mainly classified as adenocarcinomas, while bladder tumors are primarily categorized as urothelial carcinomas, with subtypes based on histological and molecular features. What is the significance of grading in the WHO classification of urinary tumors? Grading indicates the tumor's differentiation and aggressiveness, which helps guide prognosis and treatment decisions in urinary system tumors. Has the WHO classification of urinary tumors incorporated molecular and genetic features? Yes, recent updates have integrated molecular and genetic data to refine tumor classification, aiding in personalized diagnosis and therapy. Are there specific WHO classification criteria for rare urinary tumors? Yes, the WHO provides specific criteria for rare tumors such as collecting duct carcinomas, leiomyosarcomas, and other mesenchymal tumors of the urinary system. How does the WHO classification impact clinical management of urinary tumors? It provides a standardized framework for diagnosis, prognosis, and treatment planning, ensuring consistency and improved patient outcomes across different healthcare settings.

Related keywords: urinary system tumors, WHO tumor classification, urothelial carcinoma, renal cell carcinoma, bladder cancer, prostate tumors, kidney tumors, urinary tract neoplasms, tumor grading, tumor staging

Read the full article online: [WHO CLASSIFICATION OF TUMOURS OF THE URINARY SYST](#)

WHO CLASSIFICATION OF SKIN TUMOURS

WHO classification of skin tumours is an essential framework used by dermatologists, pathologists, and oncologists to categorize and understand the vast array of skin neoplasms. This classification system, developed by the World Health Organization, provides a standardized

approach to diagnosing, researching, and managing skin tumors. Accurate classification not only aids in determining prognosis but also guides appropriate treatment strategies. Skin tumors encompass a broad spectrum of benign and malignant lesions, each with distinct histopathological features and clinical behaviors. The WHO classification helps to organize this diversity into structured categories, facilitating clearer communication among healthcare professionals and enhancing patient care.

Overview of the WHO Classification of Skin Tumours

The WHO classification of skin tumours is periodically updated to incorporate new research findings, advances in diagnostic techniques, and emerging tumor entities. The current edition, often referred to as the WHO Classification of Skin Tumours (2018 edition), divides skin tumors into several major groups based on their cellular origin, histopathological characteristics, and biological behavior. These groups include carcinomas, melanocytic tumors, adnexal tumors, soft tissue tumors, and hematolymphoid tumors involving the skin.

The classification emphasizes a comprehensive approach that combines clinical features, histology, immunohistochemistry, and molecular genetics to arrive at an accurate diagnosis. It also recognizes the importance of distinguishing between benign, borderline, and malignant tumors, which has direct implications for patient management.

Major Categories of Skin Tumours in WHO Classification

1. Keratinocyte Carcinomas

Keratinocyte carcinomas are the most common malignant skin tumors, originating from the keratinocytes of the epidermis.

- **Basal Cell Carcinoma (BCC):** The most common skin cancer, characterized by nests of basaloid cells with peripheral palisading. It rarely metastasizes but can cause local destruction.
- **Squamous Cell Carcinoma (SCC):** Arises from keratinocytes of the squamous epithelium. More aggressive than BCC, with potential to metastasize, especially in immunocompromised individuals.
- **Precursor Lesions:** Includes actinic keratosis and Bowen's disease, which are considered carcinoma in situ.

2. Melanocytic Tumors

These tumors originate from melanocytes, the pigment-producing cells of the skin.

- **Benign Melanocytic Nevi:** Common moles, usually symmetrical with uniform color and well-defined borders.
- **Melanoma:** Malignant melanoma is a highly aggressive tumor with potential for metastasis. The WHO classifies melanomas based on their histological subtype, including superficial spreading, nodular, lentigo maligna, and acral melanoma.
- **Other Melanocytic Tumors:** Such as Spitz nevus and blue nevus, which can sometimes mimic melanoma.

3. Appendageal (Adnexal) Tumors

These tumors arise from the skin appendages like sweat glands, sebaceous glands, and hair follicles.

- **Eccrine and Apocrine Gland Tumors:** Includes hidradenoma, syringoma, and poroma.
- **Sebaceous Gland Tumors:** Includes sebaceous adenoma and sebaceous carcinoma.
- **Follicular Tumors:** Such as trichoepithelioma and pilomatricoma.

4. Soft Tissue Tumors

These are mesenchymal tumors originating from connective tissues within the skin.

- **Fibrous Tumors:** Such as dermatofibroma and fibrous histiocytoma.
- **Vascular Tumors:** Includes hemangiomas, Kaposi sarcoma, and angiosarcomas.
- **Lipomatous Tumors:** Such as lipomas and atypical lipomatous tumors.

5. Hematolymphoid Tumors

These involve lymphoid and hematopoietic cells infiltrating the skin.

- **Cutaneous Lymphomas:** Including mycosis fungoides and Sézary syndrome.
- **Leukemias:** Such as leukemia cutis.

Histopathological Features and Diagnostic Criteria

Accurate classification relies heavily on histopathological examination, often supplemented by immunohistochemistry and molecular studies.

Key Diagnostic Tools

- **Histology:** Microscopic examination reveals cellular morphology, growth patterns, and stromal interactions.
- **Immunohistochemistry (IHC):** Uses specific markers to distinguish tumor types, such as S100, HMB-45 for melanocytic tumors, or cytokeratins for keratinocyte tumors.
- **Molecular Genetics:** Identifies genetic mutations and chromosomal aberrations, aiding in diagnosis and targeted therapy decisions.

Importance of WHO Classification in Clinical Practice

Correctly classifying skin tumors influences treatment options, prognosis, and follow-up strategies.

Implications for Treatment

- Benign tumors may only require excision or observation.
- Malignant tumors often necessitate wide local excision, sentinel lymph node biopsy, radiotherapy, or systemic therapies.
- Some tumors, like melanoma, require staging and targeted therapy based on molecular findings.

Prognostic Significance

The classification provides prognostic information, with some tumors having a high likelihood of recurrence or metastasis, while others are essentially benign.

Emerging Trends and Future Directions

Advances in molecular pathology continue to refine skin tumor classification, leading to more personalized approaches.

Integration of Molecular Data

New genetic markers are being incorporated into the classification, allowing for better subclassification and targeted treatment options. For example, BRAF mutations in melanoma or MYC amplification in certain cutaneous lymphomas.

Development of Targeted Therapies

Understanding the molecular pathways involved in skin tumors has led to the development of targeted therapies, especially for melanoma and some adnexal tumors.

Conclusion

The WHO classification of skin tumours remains a cornerstone in dermatopathology, providing a comprehensive, standardized framework for diagnosing and managing a wide spectrum of skin neoplasms. As research advances, this classification continues to evolve, integrating new molecular insights to improve patient outcomes. Accurate classification not only enhances diagnostic precision but also facilitates personalized medicine approaches, ensuring patients receive the most effective and targeted treatments available.

This detailed overview underscores the importance of the WHO classification system in the landscape of dermatology and oncology, highlighting its role in advancing understanding, diagnosis, and management of skin tumors.

Who Classification of Skin Tumours: A Comprehensive Guide to Understanding Skin Cancer Types

When discussing skin health, one of the most critical topics is the classification of skin tumours. The who classification of skin tumours serves as a vital framework for clinicians, researchers, and patients to understand the nature, prognosis, and treatment strategies associated with various skin cancers. This structured system helps categorize tumours based on their histopathological features, origins, and biological behaviour, providing clarity in diagnosis and management.

Understanding the Importance of Skin Tumour Classification

Skin tumours encompass a wide spectrum of benign and malignant lesions, each with distinct characteristics. Proper classification influences:

- Diagnosis accuracy
- Treatment planning
- Prognostic assessment
- Research and epidemiological studies

The who classification of skin tumours offers a standardized taxonomy, facilitating uniform communication across medical disciplines and advancing scientific understanding of these lesions.

Foundation of the WHO Classification of Skin Tumours

The World Health Organization (WHO) periodically updates its classification system based on the latest research, integrating histological, immunohistochemical, and molecular data. The 2018 WHO Classification of Skin Tumours is the most recent comprehensive guide, organizing skin tumours into broad categories based on their cellular origin and behaviour.

This classification divides skin tumours primarily into:

- Epithelial tumours
- Melanocytic tumours
- Appendageal (adnexal) tumours
- Soft tissue and other mesenchymal tumours
- Hematolymphoid tumours

Within each category, tumours are further subdivided based on histopathological features.

Main Categories in the WHO Classification of Skin Tumours

- Epithelial Tumours

Epithelial tumours originate from keratinocytes or other epithelial cells of the skin. They include benign, pre-malignant, and malignant lesions.

Benign Epithelial Tumours:

- Seborrheic keratosis
- Verruca (wart)
- Sebaceous hyperplasia

Pre-malignant Epithelial Tumours:

- Actinic keratosis
- Bowen's disease (squamous cell carcinoma in situ)

Malignant Epithelial Tumours:

- Basal cell carcinoma (BCC): The most common skin cancer, arising from basal keratinocytes.
- Squamous cell carcinoma (SCC): Originates from squamous keratinocytes; potential for metastasis.
- Merkel cell carcinoma: A neuroendocrine carcinoma with aggressive behaviour.

- Melanocytic Tumours

These arise from melanocytes, the pigment-producing cells in the skin.

Benign Melanocytic Tumours:

- Melanocytic nevi (moles)
- Spitz nevi

Malignant Melanocytic Tumours:

- Malignant melanoma: The most deadly skin cancer; various subtypes include superficial spreading, nodular, lentigo maligna, and acral lentiginous melanoma.

- Appendageal (Adnexal) Tumours

Originating from skin appendages such as hair follicles, sweat glands, or sebaceous glands.

- Pilomatricoma: Tumour of hair matrix origin.
- Syringoma: Eccrine sweat duct tumour.
- Sebaceous carcinoma: Malignant tumour of sebaceous glands.
- Cylindromas and spiradenomas: Tumours of sweat gland origin.

- Soft Tissue and Mesenchymal Tumours

Includes a variety of benign and malignant tumours derived from connective tissue.

- Dermatofibroma
- Kaposi sarcoma
- Angiosarcoma

- Hematolymphoid Tumours

These are rare but include lymphomas and leukaemias that involve the skin.

- Cutaneous T-cell lymphomas (e.g., mycosis fungoides)
- Other lymphoproliferative disorders

Subclassification of Malignant Skin Tumours

Understanding the subclasses within malignant categories enhances diagnostic precision.

Basal Cell Carcinoma (BCC)

- Nodular BCC
- Superficial BCC
- Morpheaform (sclerosing) BCC
- Pigmented BCC

Features:

- Locally invasive but rarely metastasize
- Often associated with UV exposure

Squamous Cell Carcinoma (SCC)

- Well-differentiated SCC
- Poorly differentiated SCC
- Keratoacanthoma (a variant with rapid growth)

Features:

- Potential to metastasize, especially in immunocompromised patients
- Often arises in sun-exposed areas

Melanoma

- Superficial spreading melanoma
- Lentigo maligna melanoma
- Nodular melanoma
- Acral lentiginous melanoma

Features:

- Highly metastatic potential
- Requires prompt diagnosis and intervention

Molecular and Histopathological Insights

Modern WHO classification also incorporates molecular markers, such as mutations in BRAF, NRAS, and c-KIT genes, especially for melanoma and certain adnexal tumours. Histopathological examination remains the gold standard, with immunohistochemistry aiding in differentiating challenging cases.

Practical Implications of WHO Classification

The classification aids clinicians in:

- Recognizing tumour types based on clinical presentation and histology.
- Determining prognosis ? for example, melanoma subtypes have different survival rates.
- Tailoring treatment strategies, including surgical excision, radiotherapy, or systemic therapies.
- Identifying tumours requiring close follow-up due to metastatic potential.

Conclusion

The who classification of skin tumours provides an essential framework for understanding the vast array of skin neoplasms. By categorizing tumours based on their cellular origin, histological features, and molecular characteristics, healthcare professionals can improve diagnostic accuracy, inform prognosis, and optimize treatment strategies. As research advances, this classification continues to evolve, integrating new molecular data that promise to refine our understanding of skin tumour biology further. Recognizing the diversity within skin tumours underscores the importance of a multidisciplinary approach to skin cancer management, emphasizing early detection and personalized care.

Question What is the WHO classification of skin tumors based on? The WHO classification of skin tumors is based on histopathological features, cellular origin, and molecular characteristics to provide a standardized framework for diagnosis and management. How many main categories are there in the WHO classification of skin tumors? The WHO classification categorizes skin tumors into several main groups, including keratinocyte tumors, melanocytic tumors, adnexal tumors, and soft tissue tumors, among others. What are the most common keratinocyte tumors classified by WHO? The most common keratinocyte tumors classified by WHO

include basal cell carcinoma, squamous cell carcinoma, and their variants. How does the WHO classification differentiate between benign and malignant skin tumors? The classification differentiates based on histopathological criteria such as cellular atypia, invasion depth, mitotic activity, and potential for metastasis. What role does molecular pathology play in the WHO classification of skin tumors? Molecular pathology helps identify genetic mutations and markers that can refine diagnoses, predict behavior, and guide targeted therapies within the WHO classification framework. Are melanocytic tumors classified differently in the WHO system? Yes, melanocytic tumors are classified separately, including benign nevi, dysplastic nevi, and malignant melanoma, with specific subtypes based on histology and molecular features. What is the significance of the WHO classification for dermatopathologists? It provides a standardized nomenclature and diagnostic criteria, improving accuracy, facilitating research, and guiding appropriate treatment strategies. Has the WHO classification of skin tumors been updated recently? Yes, the WHO periodically updates its classifications to incorporate new scientific knowledge, with the latest edition reflecting advancements in molecular and histopathological understanding. How does the WHO classification assist in prognosis and treatment planning? By categorizing tumors based on their histological and molecular features, the classification helps predict tumor behavior and informs clinicians on optimal management approaches. What are some challenging aspects of applying the WHO classification to skin tumors? Challenges include variability in histopathological interpretation, overlapping features among tumor types, and the need for molecular testing, which may not be universally available.

Related keywords: skin tumor classification, WHO skin tumors, skin neoplasm categories, skin cancer types, dermatological tumor classification, skin tumor pathology, WHO tumor classification system, cutaneous neoplasms, skin tumor diagnosis, skin tumor epidemiology

Read the full article online: [Who Classification Of Skin Tumours](#)

WHO CLASSIFICATION OF TUMOURS OF THE LUNG PLEURA

WHO classification of tumours of the lung pleura

The World Health Organization (WHO) classification of tumours of the lung and pleura provides a comprehensive framework for diagnosing, categorizing, and understanding the diverse neoplasms that originate in these regions. This classification is pivotal for clinicians, pathologists, and researchers because it standardizes terminology, facilitates accurate diagnosis, guides treatment strategies, and aids in prognostication. This article delves into the detailed WHO classification of lung and pleural tumours, exploring the various tumour types, their histopathological features, clinical relevance, and updates from the latest WHO editions.

Overview of WHO Classification of Lung and Pleural Tumours

The WHO classification system for lung and pleural tumours was first established to address the heterogeneity of neoplasms arising within these thoracic structures. It is periodically updated to incorporate advances in pathology, molecular biology, and clinical insights. The latest WHO classification categorizes tumours into benign, borderline, and malignant entities, based on histological features, behaviour, and molecular characteristics.

The classification encompasses tumours originating from:

- Lung parenchyma (primarily epithelial tumours)
- Pleura (mesothelial tumours)
- Other rare tumours of the lung and pleura

Understanding these categories is essential for accurate diagnosis and appropriate management.

Classification of Tumours of the Lung

Lung tumours are predominantly epithelial in origin, with non-small cell lung carcinomas (NSCLCs) being the most common, followed by small cell lung carcinomas (SCLCs). The WHO classification emphasizes histological subtypes, molecular alterations, and clinical implications.

Benign Lung Tumours

Benign lung tumours are rare and usually asymptomatic. They are often discovered incidentally during imaging studies. Main benign tumours include:

- Bronchial adenomas (now termed neuroendocrine tumors of low malignant potential)
- Hamartomas
- Leiomyomas
- Hemangiomas

Malignant Lung Tumours

Malignant tumours are classified primarily based on histological features into non-small cell lung carcinomas and small cell lung carcinomas.

Non-Small Cell Lung Carcinomas (NSCLCs)

This group includes the most prevalent lung cancers and is subdivided into:

- adenocarcinoma
- squamous cell carcinoma
- large cell carcinoma

Adenocarcinoma: The most common type, especially among non-smokers. It originates from glandular epithelium and often occurs peripherally.

Squamous cell carcinoma: Typically arises centrally within the bronchi and is strongly associated with smoking.

Large cell carcinoma: A diagnosis of exclusion characterized by undifferentiated malignant cells without glandular or squamous differentiation.

Small Cell Lung Carcinoma (SCLC)

SCLC accounts for approximately 15% of lung cancers. It is characterized by small, round to oval cells with neuroendocrine features, high mitotic rate, and aggressive behavior. It often presents with extensive disease at diagnosis and responds initially to chemotherapy and radiotherapy.

Other Rare Lung Tumours

The WHO also recognizes rare tumours such as carcinoid tumours, sarcomatoid carcinomas, and lymphomas involving the lung.

Classification of Tumours of the Pleura

Pleural tumours primarily originate from mesothelial cells lining the pleura. The WHO classification emphasizes mesothelial tumours, their benign, borderline, and malignant categories.

Benign Pleural Tumours

The most common benign pleural tumour is:

- Localized fibrous tumour of the pleura (also called solitary fibrous tumour)

These are usually asymptomatic, slow-growing, and incidentally found.

Borderline and Intermediate Tumours

This category includes tumours with uncertain malignant potential, though they are less well-defined compared to malignant tumours.

Diffuse Mesothelioma of the Pleura

While classified as malignant, mesotheliomas are a distinct entity with unique pathological features.

Malignant Pleural Tumours

The most significant malignant pleural tumour is:

- Malignant mesothelioma

This tumour arises from mesothelial cells and is strongly associated with asbestos exposure.

WHO Classification of Mesothelial Tumours

Mesothelial tumours are categorized based on histological subtypes, differentiation, and cellular features:

- Epithelioid mesothelioma: The most common form, characterized by epithelial-like cells.
- Sarcomatoid mesothelioma: Composed of spindle-shaped cells resembling sarcomas.
- Biphasic (mixed) mesothelioma: Contains both epithelioid and sarcomatoid components.

Additional features include the grading of cellular atypia, mitotic activity, and invasion patterns, which influence prognosis.

Key Features and Diagnostic Criteria in WHO Classification

The WHO classification emphasizes several diagnostic criteria:

- Histopathological features: Cell morphology, growth patterns, invasion depth.

- Immunohistochemistry: Markers such as calretinin, WT-1, cytokeratin 5/6 for mesothelial origin; TTF-1, Napsin A for adenocarcinomas; p40, p63 for squamous cell carcinomas.

- Molecular features: Genetic alterations, e.g., EGFR mutations in adenocarcinoma, BRAF, KRAS mutations, and PD-L1 expression relevant for targeted therapy.

Importance of WHO Classification in Clinical Practice

The WHO classification guides clinicians by:

- Providing a standardized terminology for diagnosis and communication.
- Informing prognosis; for example, adenocarcinomas generally have better outcomes compared to small cell carcinomas.
- Influencing treatment decisions; targeted therapies depend on molecular subtypes.
- Facilitating research and epidemiological studies through consistent categorization.

Updates in the Latest WHO Classifications

The WHO periodically revises its classification to incorporate advances in pathology and molecular biology. Notable updates include:

- Recognition of new subtypes of lung adenocarcinoma (e.g., lepidic, acinar, papillary).
- Inclusion of molecular markers as diagnostic criteria.
- Clarification of borderline and pre-invasive lesions.
- Better characterization of mesothelial tumours, including epithelioid, sarcomatoid, and biphasic variants.

Conclusion

The WHO classification of tumours of the lung and pleura remains a cornerstone in thoracic pathology, fostering precise diagnosis, guiding treatment, and improving understanding of tumour biology. Its systematic approach to categorizing benign, borderline, and malignant neoplasms based on histological, immunohistochemical, and molecular features ensures consistency across institutions and enhances patient care. Staying updated with WHO revisions is essential for healthcare professionals involved in the management of thoracic tumours, ultimately leading to better outcomes for patients.

Keywords: WHO classification, lung tumours, pleural tumours, mesothelioma, lung carcinoma, benign lung tumours, malignant lung tumours, thoracic pathology, tumour histology, thoracic neoplasms

WHO Classification of Tumours of the Lung Pleura

The World Health Organization (WHO) classification of tumours of the lung and pleura represents a comprehensive framework that guides pathologists, oncologists, and researchers in diagnosing, categorizing, and managing thoracic neoplasms. This classification system, regularly updated to incorporate advances in molecular pathology and histopathology, provides a standardized language that enhances clinical decision-making and facilitates epidemiological studies. In this review, we explore the detailed taxonomy of pleural tumours as outlined by the WHO, emphasizing their histopathological features, molecular characteristics, and clinical implications.

Introduction

The pleura, a vital serous membrane enveloping the lungs and lining the thoracic cavity, is a site for a diverse array of benign and malignant tumours. Malignant pleural mesothelioma (MPM) is the most notorious primary neoplasm arising from the mesothelial cells, but secondary tumours such as metastases from lung, breast, and other carcinomas also involve the pleura. The WHO classification provides a structured approach to distinguish these entities, primarily focusing on mesothelial neoplasms, fibrous tumours, and other rare variants.

Understanding the WHO classification is essential for accurate diagnosis, prognostication, and therapeutic planning, especially given the heterogeneity of pleural tumours and the evolving landscape of molecular diagnostics.

Primary Tumours of the Pleura

Primary tumours originate from the mesothelial lining or associated stromal components. They are classified based on histopathological features, immunohistochemical profiles, and molecular

alterations.

Mesothelial Tumours

Mesothelial neoplasms are the most common primary pleural tumours and encompass benign, borderline, and malignant categories.

Benign Mesothelial Tumours

- Localized Mesothelial Hyperplasia: Reactive proliferation of mesothelial cells often associated with inflammation or trauma.
- Well-Differentiated Papillary Mesothelial Tumour (WDPMT): A rare benign entity characterized by papillary proliferation of mesothelial cells with minimal atypia and no invasion.

Malignant Mesothelial Tumours

- Malignant Mesothelioma (MM): The most significant primary malignant tumour of the pleura, with distinct histological subtypes and molecular features.

Malignant Mesothelioma Subtypes

The WHO recognizes three main histological subtypes, each with differing prognosis and therapeutic responses:

- Epithelioid Mesothelioma
- Sarcomatoid Mesothelioma
- Biphasic (Mixed) Mesothelioma

Epithelioid Mesothelioma:

Most common (~60-70%), characterized by polygonal cells forming tubules, papillae, or solid sheets. Frequently exhibits a prominent fibrous stroma and tends to have a better prognosis.

Sarcomatoid Mesothelioma:

Less common (~10-20%), composed of spindle-shaped cells resembling sarcoma. These tumours are often more aggressive and less responsive to therapy.

Biphasic Mesothelioma:

Contains both epithelioid and sarcomatoid components, with prognosis depending on the proportion of each.

Molecular and Immunohistochemical Features of Mesothelioma

- Key Immunohistochemical Markers:
 - Positive: Calretinin, WT1, D2-40 (podoplanin), Cytokeratin 5/6, Mesothelin
 - Negative: Carcinoma markers such as TTF-1, CEA, Ber-EP4
- Molecular Alterations:
 - Loss of BAP1 expression
 - CDKN2A (p16) deletion detectable via FISH
 - Rare mutations in BRAF, NRAS

Other Mesothelial Tumours

- Localized Malignant Mesothelioma: Rare, similar to diffuse but confined to localized areas.
- Multicystic Mesothelioma: Generally benign, presenting as multicystic lesions.

Fibrous and Other Tumours of the Pleura

Aside from mesothelial origins, the pleura can host fibrous and miscellaneous tumours.

Solitary Fibrous Tumour (SFT)

- Originates from fibroblastic mesenchymal cells.
- Usually benign but can be malignant.
- Features: Well-circumscribed, spindle cells with high vascularity, "patternless" pattern, and CD34 positivity.
- Malignant SFTs show increased cellularity, atypia, mitoses, necrosis.
- Molecular hallmark: NAB2-STAT6 gene fusion detectable via immunohistochemistry (STAT6

nuclear positivity).

Fibrosarcoma and Other Sarcomas

Rare; includes fibrosarcoma, malignant peripheral nerve sheath tumour, and angiosarcoma. These are generally distinguished through histomorphology and immunoprofile.

Mesenchymal Tumours with Other Differentiations

- Lipomas and Liposarcomas
- Hemangiopericytoma

Secondary Tumours of the Pleura

Metastatic involvement of the pleura is common, especially from lung, breast, and other carcinomas.

Common Metastases

- Lung Carcinomas: Adenocarcinoma, squamous cell carcinoma
- Breast Carcinoma
- Gastrointestinal Tumours
- Lymphomas and Leukemias

Immunohistochemistry and molecular studies help determine the primary site, especially in poorly differentiated tumours.

Rare and Unusual Pleural Tumours

- Mesenchymal chondrosarcoma
- Malignant fibrous histiocytoma
- Synovial sarcoma

These are exceedingly rare but are recognized within the WHO framework based on histopathology and molecular features.

Diagnostic Challenges and Advances

Accurate classification relies on a combination of histopathology, immunohistochemistry, and

increasingly, molecular diagnostics.

Role of Molecular Diagnostics

- Detection of BAP1 loss and CDKN2A deletions for mesothelioma
- Identification of NAB2-STAT6 fusion in SFT
- Next-generation sequencing panels for rare entities

Diagnostic Algorithms

- Use of panel immunohistochemistry to differentiate mesothelioma from metastatic carcinoma
- Employing molecular markers for ambiguous cases

Clinical Implications of WHO Classification

The WHO classification informs prognosis and guides management strategies:

- Benign vs. Malignant: Surgery for benign tumours; multimodal therapy for mesothelioma.
- Histological Subtypes: Epithelioid mesotheliomas have better outcomes.
- Molecular Features: Potential targets for novel therapies.

Conclusion

The WHO classification of tumours of the lung and pleura offers a detailed and nuanced taxonomy that reflects advances in our understanding of thoracic neoplasms. It emphasizes the importance of integrating histopathological, immunohistochemical, and molecular data to achieve accurate diagnosis. As research continues to evolve, especially in molecular genetics and targeted therapies, this classification will likely undergo further refinements, ultimately improving patient outcomes through tailored treatment approaches.

Understanding the diverse spectrum of pleural tumours aids clinicians and pathologists in making precise diagnoses, prognostic assessments, and guiding appropriate management. The ongoing evolution of the WHO classification underscores the importance of multidisciplinary collaboration in thoracic oncology.

QuestionAnswer What is the WHO classification of tumors of the lung and pleura? The WHO classification of tumors of the lung and pleura is a standardized system that categorizes benign and malignant tumors based on histological features, genetic profiles, and clinical behavior to guide

diagnosis and treatment. How are mesotheliomas classified in the WHO system? In the WHO classification, mesotheliomas are categorized into epithelioid, sarcomatoid, biphasic (mixed), and rare variants, based on their cellular morphology and immunohistochemical profiles. What are the main types of primary lung carcinomas according to WHO? The main types include non-small cell lung carcinomas (adenocarcinoma, squamous cell carcinoma, large cell carcinoma) and small cell lung carcinoma, each with distinct histological and molecular features. How does the WHO classify benign tumors of the lung and pleura? Benign tumors are classified as hamartomas, papillomas, lipomas, and other rare benign neoplasms, characterized by their benign histological appearance and non-invasive behavior. What is the significance of the WHO classification in clinical management? The WHO classification helps in accurate diagnosis, prognosis, and selection of appropriate treatment strategies by providing a standardized framework for tumor categorization. Are genetic features incorporated into the WHO classification of lung and pleural tumors? Yes, the WHO classification increasingly incorporates molecular and genetic markers that aid in subclassification, prognosis, and targeted therapy options. How are metastatic tumors to the lung and pleura classified in the WHO system? Metastatic tumors are classified based on their primary origin and histological similarity, with common types including metastases from breast, colon, and other cancers, but they are not part of the primary WHO tumor classification. What updates have recent WHO classifications introduced for lung and pleural tumors? Recent updates include enhanced molecular characterization, recognition of new tumor variants, and refined criteria for differentiating benign from malignant lesions to improve diagnostic accuracy. How does the WHO classification address rare tumors of the lung and pleura? The WHO classification provides specific categories for rare tumors such as solitary fibrous tumors, carcinoids, and other uncommon neoplasms, facilitating their recognition and management. Why is the WHO classification important for pathologists and clinicians? It ensures consistent diagnosis, informs prognosis, guides treatment decisions, and fosters research by providing a comprehensive, evidence-based framework for tumor categorization.

Related keywords: lung tumors, pleural tumors, WHO classification, thoracic oncology, mesothelioma, lung cancer types, pleural mesothelioma, malignant pleural effusion, tumor histology, lung neoplasms

Read the full article online: [who classification of tumours of the lung pleura](#)

who classification of tumours of endocrine organs

WHO classification of tumours of endocrine organs is a comprehensive system developed by the World Health Organization to categorize and diagnose tumors originating from endocrine glands. This classification provides a standardized framework that aids pathologists, endocrinologists, and

oncologists in understanding the nature, prognosis, and appropriate treatment strategies for these tumors. Given the complexity and diversity of endocrine tumors, the WHO classification has evolved over the years to incorporate advances in molecular pathology, histopathology, and clinical research.

Introduction to Endocrine Tumors and Their Significance

Endocrine tumors are neoplasms arising from endocrine glands such as the thyroid, parathyroid, adrenal glands, pituitary, and neuroendocrine cells scattered throughout various organs. These tumors can be benign or malignant and often produce hormones, leading to diverse clinical syndromes. Accurate classification is vital for prognosis determination, therapeutic decision-making, and research.

Historical Development of WHO Classification of Endocrine Tumors

The WHO classification was first introduced in 2004 and has undergone multiple updates, notably in 2017. These revisions reflect new insights from molecular genetics, immunohistochemistry, and clinical behavior patterns. The current system emphasizes a combined approach that considers morphology, immunophenotype, and molecular features, facilitating precise diagnosis and personalized medicine.

Major Categories in WHO Classification of Endocrine Tumours

The WHO classification divides endocrine tumors primarily based on the organ of origin and tumor behavior. The main categories include:

- Thyroid tumors
- Adrenal tumors
- Parathyroid tumors
- Pituitary tumors
- Neuroendocrine tumors of other organs

Each category comprises benign, uncertain malignant potential, and malignant tumors, with specific subtypes and grading systems.

Thyroid Tumours

Thyroid tumors are among the most common endocrine neoplasms. They are classified into several categories:

Differentiated Thyroid Carcinomas

These tumors retain some features of normal thyroid tissue and are generally associated with a better prognosis.

- **Papillary Thyroid Carcinoma (PTC):** The most common type, characterized by papillary structures and nuclear features such as clearing and grooves.
- **Follicular Thyroid Carcinoma (FTC):** Comprises follicular structures; invasion through the capsule or blood vessels signifies malignancy.
- **Hürthle Cell Carcinoma:** A variant of FTC with oncocytic features.

Medullary Thyroid Carcinoma (MTC)

Arises from parafollicular C cells and secretes calcitonin. It has familial and sporadic forms and is associated with RET mutations.

Anaplastic Thyroid Carcinoma

A highly aggressive undifferentiated carcinoma with poor prognosis.

Other Thyroid Tumors

Includes tumors such as diffuse sclerosing carcinoma and benign adenomas.

Adrenal Tumours

Adrenal tumors originate from the cortex or medulla, with distinct classifications.

Adrenocortical Tumors

These include benign adenomas and malignant carcinomas.

- **Adrenocortical Carcinoma (ACC):** Malignant, often secreting hormones like cortisol, and associated with poor prognosis.
- **Adrenocortical Adenoma:** Usually benign and hormonally active or inactive.

Pheochromocytoma and Paragangliomas

Arise from chromaffin cells in the adrenal medulla or paraganglia.

- **Pheochromocytoma:** Located in adrenal medulla; secretes catecholamines.
- **Paranglioma:** Extra-adrenal; can be sympathetic or parasympathetic.

Other Adrenal Tumors

Rare tumors include myelolipomas and metastases.

Parathyroid Tumours

The most common parathyroid tumor is the parathyroid adenoma, which causes primary hyperparathyroidism.

Benign Parathyroid Tumors

- Parathyroid adenoma: The majority of cases.

Parathyroid Carcinoma

A rare but aggressive malignant tumor requiring surgical excision.

Pituitary Tumours

Pituitary tumors are classified based on size, hormone secretion, and histology.

Adenomas

Most common, benign tumors classified by the hormone they produce:

- Prolactinomas
- Somatotroph adenomas (growth hormone-secreting)
- Corticotroph adenomas (ACTH-secreting)
- Other hormone-secreting adenomas

Pituitary Carcinomas

Extremely rare malignant tumors with metastatic potential.

Others

Include Rathke cleft cysts and craniopharyngiomas, which are non-neoplastic lesions.

Neuroendocrine Tumours of Other Organs

Neuroendocrine tumors (NETs) are a diverse group originating from neuroendocrine cells in various tissues, including the pancreas, lungs, and gastrointestinal tract.

Pancreatic Neuroendocrine Tumours (PanNETs)

Classified based on differentiation, hormonal activity, and grading.

- Well-differentiated neuroendocrine tumors (NETs)
- Neuroendocrine carcinomas (NECs)

Gastroenteropancreatic Neuroendocrine Tumours (GEP-NETs)

Includes tumors of the stomach, small intestine, colon, and rectum.

Lung Neuroendocrine Tumors

Divided into typical carcinoids, atypical carcinoids, large cell neuroendocrine carcinomas, and small cell lung carcinomas.

Grading and Staging in WHO Classification

The WHO system incorporates grading based on mitotic activity, Ki-67 proliferation index, and histological features to predict tumor behavior. Staging follows the TNM system, considering tumor size, lymph node involvement, and metastasis.

Molecular Pathology and Its Role in Classification

Recent updates emphasize molecular alterations, such as mutations in RET, RAS, TP53, and others, which refine diagnosis and enable targeted therapy. For example:

- RET mutations are prominent in medullary thyroid carcinoma.
- BRAF mutations are common in papillary thyroid carcinoma.
- MDM2 amplification may suggest certain adrenal tumors.

Molecular testing complements histopathology, especially in ambiguous cases.

Clinical Implications of WHO Classification

Accurate classification influences treatment choices, prognosis, and follow-up strategies. For example:

- Benign tumors may require surgical excision with curative intent.
- Malignant tumors often need multimodal therapy, including surgery, radiotherapy, and targeted agents.
- Understanding tumor subtype informs prognosis and genetic counseling, especially in familial syndromes like MEN (Multiple Endocrine Neoplasia).

Conclusion

The WHO classification of tumours of endocrine organs provides a vital framework that integrates morphological, immunohistochemical, and molecular features to deliver precise diagnoses. As research advances, this classification continues to evolve, reflecting our growing understanding of the pathogenesis and behavior of endocrine tumors. Clinicians and pathologists must stay updated with these revisions to optimize patient care, facilitate research, and improve outcomes for individuals affected by these diverse neoplasms.

Keywords: WHO classification, endocrine tumors, thyroid neoplasms, adrenal tumors, parathyroid tumors, pituitary tumors, neuroendocrine tumors, tumor grading, molecular pathology

WHO classification of tumours of endocrine organs is an essential framework that provides a standardized approach to diagnosing, categorizing, and understanding tumors arising from endocrine tissues. These classifications are integral for pathologists, endocrinologists, oncologists, and researchers alike, as they guide clinical management, prognostication, and research directions. Since the World Health Organization's (WHO) first publication on this topic, the classification system has undergone multiple updates to incorporate advances in molecular biology, histopathology, and clinical outcomes. This review aims to provide a comprehensive overview of the WHO classification of endocrine tumors, discussing its structure, key categories, updates, strengths, limitations, and clinical implications.

Introduction to WHO Classification of Endocrine Tumours

The WHO classification of tumours of endocrine organs is a globally recognized system that categorizes tumors based on histological features, immunohistochemical profiles, molecular genetics, and clinical behavior. It serves as an authoritative reference for tumor diagnosis and classification, promoting consistency and comparability across different institutions and studies. The latest edition, published in 2022, reflects significant advances in understanding the molecular

underpinnings of endocrine tumors, which have led to refinements in classification criteria.

The classification encompasses tumors originating from various endocrine organs, including the thyroid, parathyroid, adrenal glands, pituitary, neuroendocrine neoplasms of the pancreas, and other neuroendocrine tumors (NETs). It distinguishes benign from malignant lesions, identifies variants with distinct behavior, and integrates molecular insights for a more precise categorization.

Structure of the WHO Classification System

The WHO classification is organized by organ system, with each chapter dedicated to a specific endocrine organ. Within each chapter, tumors are classified into categories based on their histogenesis, differentiation, and aggressiveness. The key elements include:

- Benign tumors: e.g., adenomas
- Borderline or uncertain malignant potential: e.g., certain tumors with atypical features
- Malignant tumors: e.g., carcinomas, neuroendocrine carcinomas

The classification also incorporates:

- Grading: based on proliferative activity
- Staging: for clinical prognosis
- Molecular features: genetic mutations associated with tumor types

This systematic approach enhances diagnostic accuracy and informs prognosis and management strategies.

Thyroid Tumours

The thyroid gland harbors a variety of neoplasms, with papillary thyroid carcinoma (PTC) being the most common.

Benign Thyroid Tumours

- Follicular adenoma: encapsulated, benign tumor of follicular cells.
- Features:
 - Well-circumscribed, non-invasive
 - No capsular or vascular invasion

Thyroid Carcinomas

- Papillary thyroid carcinoma (PTC):
- Most common thyroid malignancy
- Characterized by papillary structures, nuclear features like grooves and inclusions
- Molecular hallmark: BRAF mutations, RET/PTC rearrangements
- Follicular thyroid carcinoma:
- Invades capsule or blood vessels
- Molecular features: RAS mutations, PAX8-PPAR γ rearrangements
- Medullary thyroid carcinoma (MTC):
- Arises from parafollicular C cells
- Associated with RET mutations, MEN syndromes
- Anaplastic thyroid carcinoma:
- Highly aggressive, undifferentiated tumor
- Poor prognosis

Features & Pros/Cons:

Feature	Pros	Cons
Molecular profiling	Guides targeted therapies	Not always available in resource-limited settings
Histological classification	Standardized diagnosis	Overlap in some subtypes complicates diagnosis

Parathyroid Tumours

Most parathyroid tumors are benign adenomas responsible for primary hyperparathyroidism.

Benign Parathyroid Tumours

- Parathyroid adenoma:
- Encapsulated, monoclonal proliferation
- Features: Chief cell proliferation, sometimes with oxyphil cells
- Features & Considerations:
- Usually solitary
- Can be distinguished from hyperplasia by histology

Malignant Parathyroid Tumours

- Very rare
- Parathyroid carcinoma:
- Invasive growth, potential metastasis
- Features: Trabecular growth, capsular invasion, mitoses

Features & Pros/Cons:

| Feature | Pros | Cons |

|---|---|---|

| Histology | Helps differentiate benign from malignant | Limited by overlapping features and rarity of carcinomas |

Adrenal Gland Tumours

Adrenal tumors include benign adenomas, malignant carcinomas, and neuroendocrine tumors.

Adrenal Cortical Tumours

- Adenomas: benign, non-functional or functional
- Carcinomas: aggressive, associated with hormonal syndromes
- WHO criteria: Weiss score assesses malignancy potential

Pheochromocytomas and Paragangliomas

- Neuroendocrine tumors arising from chromaffin cells
- Features:
- Pheochromocytomas are adrenal
- Paragangliomas are extra-adrenal
- Genetic mutations: SDHx, RET, VHL, NF1

Features & Pros/Cons:

| Feature | Pros | Cons |

|---|---|---|

| Molecular genetics | Allows targeted therapy and risk stratification | Requires advanced testing infrastructure |

Pituitary Tumours

Most are benign adenomas, with a few aggressive or malignant variants.

Types of Pituitary Tumours

- Lactotroph adenomas: prolactin-secreting
- Somatotroph adenomas: growth hormone-secreting
- Corticotroph adenomas: ACTH-secreting, causing Cushing's disease
- Non-functioning adenomas: do not secrete hormones

Features & Pros/Cons:

| Feature | Pros | Cons |

|---|---|---|

| Immunohistochemistry | Helps identify tumor subtype | Hormone levels often guide diagnosis more directly |

Neuroendocrine Tumours of the Pancreas and Other Sites

This includes pancreatic neuroendocrine tumors (PNETs) and other neuroendocrine neoplasms.

Classification of PNETs

- Well-differentiated tumors:
 - Functioning (insulinoma, gastrinoma, etc.)
 - Non-functioning
 - Graded based on proliferative index (Ki-67) and mitotic rate
- Poorly differentiated neuroendocrine carcinomas:
 - High-grade, aggressive tumors
 - Features: high mitotic rate, necrosis

Features & Pros/Cons:

| Feature | Pros | Cons |

|---|---|---|

| WHO grading | Provides prognostic information | Requires precise proliferation assessment |

Advantages of the WHO Classification

- Standardization: Ensures consistent terminology across institutions.
- Integration of Molecular Data: Reflects current understanding of tumor genetics.
- Prognostication: Helps predict clinical behavior based on histology and molecular features.
- Guides Treatment: Facilitates tailored therapeutic approaches, including targeted therapies.
- Research Facilitation: Provides a framework for clinical and translational research.

Limitations and Challenges

- Complexity: The detailed molecular classifications can be resource-intensive.
- Variability in Access: Not all centers have access to advanced molecular diagnostics.
- Evolving Nature: Rapid advances mean classifications need frequent updates.
- Overlap in Features: Some tumors display overlapping histological and molecular features, complicating diagnosis.
- Prognostic Uncertainty: Some tumors with similar histology behave differently clinically.

Clinical Implications of WHO Classification

The WHO classification influences various aspects of patient care:

- Diagnosis: Accurate tumor typing determines prognosis and management.
- Treatment Planning: Molecular insights enable targeted therapy and personalized medicine.
- Follow-Up: Grading and staging inform surveillance strategies.
- Research and Trials: Uniform classification allows for better patient stratification in clinical trials.

Conclusion

The WHO classification of tumours of endocrine organs represents a cornerstone in endocrine pathology, integrating histopathological features with molecular genetics to refine tumor diagnosis,

prognostication, and treatment. While it offers numerous advantages, ongoing updates and integration of novel molecular techniques are necessary to overcome current limitations. As our understanding of tumor biology deepens, future iterations will likely become even more precise, facilitating personalized approaches that improve patient outcomes. Ultimately, the WHO classification remains a dynamic and vital tool in the management of endocrine neoplasms, reflecting the evolving landscape of tumor biology and clinical practice.

QuestionAnswer What is the WHO classification of tumors of endocrine organs based on? The WHO classification of tumors of endocrine organs is based on histopathological features, including cell type, differentiation, molecular characteristics, and tumor behavior, to provide a standardized framework for diagnosis and prognosis. How are neuroendocrine tumors of the pancreas categorized in the WHO classification? In the WHO classification, pancreatic neuroendocrine tumors are categorized into well-differentiated neuroendocrine tumors (NETs) grades 1 and 2, poorly differentiated neuroendocrine carcinomas (NECs), and mixed neuroendocrine-non-neuroendocrine neoplasms (MiNENs), based on differentiation and proliferative activity. What are the key features used to differentiate benign from malignant endocrine tumors in WHO classification? Key features include tumor size, invasion into surrounding tissues, mitotic rate, Ki-67 proliferation index, presence of metastasis, and cellular atypia, which help determine whether an endocrine tumor is benign or malignant. How does the WHO classify medullary thyroid carcinoma? The WHO classifies medullary thyroid carcinoma as a malignant tumor arising from parafollicular C cells, characterized by amyloid stroma and calcitonin production, and distinguishes it from other thyroid neoplasms based on histology and immunohistochemistry. What is the significance of molecular markers in the WHO classification of endocrine tumors? Molecular markers such as MEN1, DAXX, ATRX, and RET mutations are increasingly incorporated into the WHO classification to improve diagnostic accuracy, predict behavior, and guide targeted therapies for endocrine tumors.

Related keywords: endocrine tumors, WHO tumor classification, endocrine neoplasms, endocrine gland tumors, neuroendocrine tumors, endocrine pathology, tumor grading, tumor staging, endocrine tumor types, tumor histology

Read the full article online: [Who Classification Of Tumours Of Endocrine Organs](#)